

References bibliographiques de la campagne PEL 2001 :

- 1) Bergeron J.P. 1997. Effect of strong winds on the nutritional condition of anchovy (*Engraulis encrasicolus* L.) larvae in the Bay of Biscay, N.E. Atlantic, as inferred from an early field application of the DNA/C index. ICES Symposium on recruitment dynamics - Baltimore 22-24/9/1997.
- 2) Checkley, D. ; Ortner, P. ; Settle, L. ; Cummings, S., 1997, A continuous underway fish egg sampler, Fisheris Oceanography, 6(2) : 58-73.
- 3) Herblant Alain, Daniel Delmas, Pierre Laborde, Benoit Sautour, Felipe Artigas., 1998. Phytoplankton spring bloom of the Gironde plume waters in the Bay of Biscay: early phosphorus limitation and food-web consequences. *Oceano. Acta*, vol. 21, 279-291.
- 4) Lazure P., Jegou A.M. (1998): 3D modelling of seasonal evolution of Loire and Gironde plumes on Biscay Bay Continental Shelf. In: "5eme Colloque International d'Oceanographie du Golfe de Gascogne". La Rochelle, Avril 1996. *Oceanologica Acta* , 21, 2, pp165-177.
- 5) Masse J., C. Koutsikopoulos and W. Patty, 1996 - The structure and spatial distribution of pelagic fish schools in multi-species clusters : an acoustic study. *ICES Journal of Marine Science*, 53: 155-160.