

REFERENCES BIBLIOGRAPHIQUES :

- Ahagon, N., Tanaka, Y and Ujiiie, H (1993) Florisphaera Profunda, a possible nannoplankton indicator of late Quaternary changes in sea water turbidity at the northwestern margin of the Pacific. *Marine Micropaleontology*, 22 : 255-273
- Bassinot F C, Melieres F, Levi C, Ghelen M., Labeyrie L., and H. Elderfield (2001). Improving Mg/Ca paleothermometer : Correction of dissolution effects on Mg/Ca ratio of foraminifera shells. 7th International Conference on Paleoceanography, September 16-22, 2001, Sapporo, Japan. Abstract : 79
- Bassinot, F., Levi, C., Melières, F., Ghelen, M., Labeyrie, L., and Elderfield, H. Improving the Mg/Ca paleothermometry : Correction of dissolution effects on Mg/Ca of planktonic foraminifera based on calcite crystallinity index, example of *Globigerinoides ruber*. Soumis a G-cubed
- Beaufort L.(1996) Dynamics of the monsoon in the equatorial Indian Ocean over the last 260000 years. *Quaternary International*, 31 : 13-18
- Oda, M. and Takemoto, A.(1992). Planktonic foraminifera and paleoceanography in the domain of the Kuroshio Current around Japan during the last 20,000 years. *The Quaternary Research*, 31: 341-357.
- Ujiie H.& Ujiie Y. (1999). Late Quaternary course changes of the Kuroshio current in the Ryukyu Arc region, northwestern Pacific Ocean. *Marine Micropaleontology*, 37 : 23-40.
- Xu, X., and Oda, M. (1999). Surface-water evolution of the eastern East China Sea during the last 36,000 years. *Marine Geology*, 156: 285-304