

REFERENCES BIBLIOGRAPHIQUES

- DESAUNAY Y., GUERAULT D., LE PAPE O. ET POULARD J.-C. 2006. Changes in occurrence and abundance of northern / southern flatfishes over a 20-year period in a coastal nursery area (Bay of Vilaine) and on the eastern continental shelf of the Bay of Biscay. *Scientia Marina* 70: 193-200.
- GILLIERS C., AMARA R., BERGERON J.P. ET LE PAPE O. 2002. Quality of flatfish nursery habitats in the Eastern Channel and Southern Bight of the North Sea. Communication. Fifth International Symposium on Flatfish Ecology, 3-7 November 2002, Port Erin Marine Laboratory, Isle of Man, UK.
- GILLIERS C., AMARA R., BERGERON J.P., LE PAPE O. ET DESAUNAY Y. 2002. Comparison of Flatfish nursery habitats in the Bay of Biscay using growth and condition. Communication. 8ème colloque international d'océanographie du golfe de Gascogne, 10-12 Avril 2002, IEO, Gijon, SP.
- LE BRIS H. ET GLEMAREC M. 1995. Macrobenthic communities of an oxygen of undersaturated coastal ecosystems: The Bay of Vilaine (southern Brittany). *Oceanologica acta*. 18: 573-581.
- LE BRIS H. 1988. Fonctionnement des écosystèmes benthiques côtiers au contact d'estuaires : la rade de Lorient et la baie de Vilaine. Thèse d'Océanologie biologique. Université de Bretagne Occidentale, Brest, 311 pages
- LE PAPE O., BAULIER L., CLOAREC A., MARTIN J., LE LOCH F. ET DESAUNAY Y. 2007. Habitat suitability for the juvenile common sole (*Solea solea*, L.) in the Bay of Biscay (France) : a quantitative description using indicators based on benthic fauna. *Journal of Sea Research* 57 : 126-136.
- LE PAPE O., CHAUVET F., DESAUNAY Y. ET GUERAULT D. 2003a. Relationship between interannual variations of the river plume and the extent of nursery grounds for the common sole (*Solea solea*, L.) in Vilaine Bay. Effects on recruitment variability. *Journal of Sea Research*, 50 : 177-185.
- LE PAPE O., CHAUVET F., MAHEVAS S., LAZURE P., GUERAULT D. ET DESAUNAY Y. 2003b. Quantitative description of habitat suitability for the juvenile common sole (*Solea solea*, L.) and contribution of different habitats to the adult population in the Bay of Biscay (France). *Journal of Sea Research*, 50 : 139-149.
- LE PAPE O., DESAUNAY Y. & GUERAULT D. 2003c. Relationship between fluvial discharge and sole (*Solea solea* L.) recruitment in the Bay of Biscay (France). ICES Marine Symposia, 219 : 241-248.
- LE PAPE O., CHAUVET F., MAHEVAS S., LAZURE P., GUERAULT D. ET DESAUNAY Y. 2002. Rôle des forçages naturels et anthropiques sur les écosystèmes et les principales ressources halieutiques du Golfe de Gascogne. In Rapport à mi-parcours du Programme National d' Ecologie Côtière, Chantier Golfe de Gascogne. A. Herblard ed. Ifremer: 94-98.
- NICOLAS D., LE LOC'H F., DESAUNAY Y., BLANCHET A.,HAMON D., LE PAPE O. (accepté). Relationships between benthic macrofauna and habitat suitability for juvenile common sole (*Solea solea*, L.) in the Vilaine estuary (Bay of Biscay, France) nursery ground. *Estuarine, coastal and shelf science*.

- POULARD J.-C., DESAUNAY Y., GUERAULT D., LE PAPE O. ET MARTIN J. 2002. Spatial variation in demersal assemblages and benthic macrofauna on coastal nursery grounds in the Bay of Biscay. Ifremer. Communication au 8ème colloque international d'océanographie du golfe de Gascogne. Gijón, 10-12 avril 2002.
- ROCHET M.-J., TRENKEL V, BELLAIL R, COPPIN F, LE PAPE O, MAHE JC, MORIN J, POULARD JC, SCHLAICH I, SOUPLET A, VERIN Y, BERTRAND J. 2005. Combining indicator trends to assess ongoing changes in exploited fish communities: diagnostic of communities off the coasts of France. ICES Journal of Marine Science 62: 1647-1664.
- TRENKEL V.M. AND ROCHET M.-J . 2003. Performance of indicators derived from abundance estimates for detecting the impact of fishing on a fish community. Canadian Journal of Fish and Aquatic Sciences 60: 67–85.